


ESRF | The European Synchrotron

BM29 NEWS



microfluidics: crystallisation in situ;

complementarity with SANS: joint access

2D

detector

2) BM29 OUTLOOK

KEY FEATURES

- beamline is dedicated exclusively to small-angle scattering experiments of biological <u>macromolecules in solution</u>;
- minimal sample requirements: 10 μL with c << 1 mg/mL;
- high throughput facility "easy" to use in typical data collection times
- a) sample changer mode 10 s (cycle loading-exposure-cleaning ~ 1.5 min)
- b) **HPLC on-line** mode 30 min (column elution time);
- providing <u>automated</u> data analysis and data tracking;
- access through <u>Rolling procedure</u> or BAG system: proposals any time and beam time in a few weeks (if successful and time available);
- <u>SANSAXS BAG</u>: one trip to Grenoble for ILL and ESRF experiments on same samples: SAS with neutrons and X-rays.


BM29 OUTLOOK – AUTOMATISATION

ROBOT/HPLC




EDNA


EXPERIMENTAL parameters



BsxCuBE




ISPyBB sample tracking and data flow






BM29 OUTLOOK = SAMPLE CHANGER + HPLC SYSTEM



ION EXCHANGE CROMATOGRAPHY




Separation based on surface charge









Salt concentration (linear gradient)

Advantage:

- sample can be loaded at low concentrations (avoids sample loss due to aggregation
- relatively high flow rate (mL/min) reduces the risk of radiation damage

Disadvantage:

- buffer composition intrinsically unknown


Collaboration:
Martha Breninich (EMBL,
Grenoble)
Stephanie Hutin (IBS,
Grenoble)

Figures: GE Healthcare




IEC-SAXS online: Small-POX HELICASE-PRIMASE d5

salt concentration (step gradient)









• IEC can separate similarly sized macromolecules

using careful background correction,
 high quality SAXS data can be obtained

"Online ion-exchange chromatography for small-angle X-ray scattering"
S. Hutin, M. Brennich, B. Maillot and A. Round Acta Cryst. (2016) D72, 1090–1099.





MICROFLUIDICS INTEGRATION AT BM29



COUPLING DIGITAL MICROFLUIDICS AND SAXS AT BM29

External triggering when coupling digital microfluidics and SAXS:






CRYSTALLISATION IN SITU

surfactant Each droplet acts as a microreactor in which the concentration 1000 pH, ligands, and additives) can be fine-tuned, wit As the conditions can be adjusted dynamically by 800 range of conditions can be investigated quickly a crystals Glucose isomerase + surfactant 200 crystallisation agent buffer protein a (nm **FLOW** oil with surfactant 200 μm

JOINT SANSAXS EXPERIMENT




% D₂O in H₂O/D₂O solvent

SANS contrast variation experiments obtained by exchanging the solvent for deuterated or partially deuterated solvent enhances the signal from one component of a complex.

40% D₂O buffer

0% D₂O buffer



D22 at ILL

20




Joint access – ESRF and ILL SAS experiments during one trip to Grenoble.

SANSAXS BAG from June 2016: when SAN/XS proposal accepted (and scheduled) by ILL → beamtime (typically 1 shift) on BM29 within a few days of ILL experiment.

100

100% D₂O buffer

JOINT SANSAXS STATS







BM29 NEWS: NEW SAMPLE PREPARATION LAB



BM29 NEWS: NEW SAMPLE PREPARATION LAB



BM29 NEWS: NEW SAMPLE PREPARATION LAB







https://exi.esrf.fr/saxs


http://ispyb.esrf.fr

- new GUI, which separates database and user interface;
- will replace ISPyB, new features added only to EXI;
- in development, but in use already, accessible from outside;
- much quicker access, personalized accounts, searching by key word, LC, date



EXTENDED ISPYB = EXI







EXTENDED ISPYB = EXI

