

# Microbeam radiation therapy (MRT): Milestones - Clinical prospects

J.A. Laissie<sup>1</sup>, H. Blattmann<sup>1</sup>, M.A. Grotzer<sup>2</sup>, B. Kaser-Hotz<sup>3</sup>, D.N. Slatkin<sup>4</sup>,  
H.P. Wagner<sup>5</sup>

<sup>1</sup> Institut für Pathologie, Universität Bern, CH-3010 Bern, Switzerland

<sup>2</sup> Universitäts-Kinderkliniken, CH-8032 Zürich, Switzerland

<sup>3</sup> Vetsuisse Fakultät, Universität Zürich, CH-8057 Zürich, Switzerland

<sup>4</sup> Brookhaven National Laboratory, Upton, New York 11973-5000, USA

<sup>5</sup> Universitäts-Kinderkliniken des Inselspitals, CH-3010 Bern, Switzerland

**Key words:** synchrotron X-ray microbeam radiotherapy - MRT - radiosurgery - pediatric neuro-oncology - CNS tumors

## Rationale and objectives

Collateral damage to vital normal tissues during radiotherapy can be reduced by using three-dimensional treatment planning and external sources of ionizing radiation. Nevertheless, pediatric oncologists try to postpone or forgo any radiosurgery or radiotherapy, especially for children under three years old because irradiating a child's CNS entails a substantial risk of dysfunctional central nervous system (CNS) development<sup>1, 2, 3</sup>. In radiosurgery, spatially accurate and highly conformal beams of radiation are targeted toward a well-delineated tumor in a single session<sup>5</sup>. High-dose radiosurgery using multiple millimeters-wide beams of X rays was first described in 1909<sup>6</sup>. In modern radiosurgery<sup>7</sup>, multiple millimetres-wide beams of linac-generated X rays, or of gamma rays, converge in the target. Might MRT, a radiosurgery mediated by multiple microscopically thin planar beams of synchrotron-generated X rays, yield larger therapeutic indices for CNS tumors than other forms of radiosurgery or radiotherapy?

## Methods

MRT, a spatially fractionated radiotherapy, uses an array of microscopically thin, nearly parallel synchrotron-generated X-ray beams<sup>8, 9</sup>. Peak radiation doses are up to fifty times higher than in other radiosurgeries. Unlike conventional radiotherapy, for which the effect of changing an irradiation parameter, e.g., the dose fractionation schedule, is predictable, methods to predict the effect of varying an MRT parameter are only beginning to be developed. Among MRT parameters are array width and height, slit width, spacing of microbeams, energy spectrum, changes in tissue dose microdistribution with tissue depth and, possibly in the future, the schedule selected for temporal fractionation of multidirectional MRT.

## Results

In animal experiments, MRT has shown unprecedented sparing of normal radiosensitive tissues as well as preferential damage to malignant tumor tissues growing into and around such normal tissues in laboratory animals<sup>10, 11, 12-15, 16-18, 19, 21, 22, 23-25, 26-28</sup>. MRT research at the National Synchrotron Light Source (NSLS), Upton, New York, and at the European Synchrotron Radiation Facility (ESRF), Grenoble, France, has shown that single-fraction, unidirectional MRT yields a larger therapeutic index than does single-fraction unidirectional broad beam irradiation for the intracerebral rat 9L gliosarcoma (9L GS)<sup>13, 15-17, 23-26</sup> and for the transplanted subcutaneous murine mammary carcinoma

EMT-613-15, as does bidirectional (orthogonally cross-fired) MRT for the subcutaneously transplanted, aggressively invasive, extraordinarily radioresistant murine squamous cell carcinoma VII20.

Since postponing radiotherapy may jeopardize survival of some children with brain tumors<sup>29</sup>, MRT has been undergoing and undergoes experimental assessment in living animals because it is believed to be potentially useful for inhibiting children's brain tumors while sparing nearby normal CNS tissues, which should reduce the burden of malignant cells and, therefore, enhance the effectiveness of ancillary therapies<sup>30</sup>. The relative sparing by X-ray microbeams of normal tissues of vertebrates - particularly of their normal central nervous system tissues - has been documented in suckling and adult rats<sup>18, 24, 26, 28</sup>, duck embryos<sup>12</sup>, and weanling piglets<sup>19</sup>. These preclinical results, although encouraging, are not yet sufficient to justify a Phase I (safety) trial of MRT for human patients because they are all based on small animal models, except for a set of studies at the ESRF that used the normal piglet cerebellum<sup>16</sup>. All other normal-tissue microbeam-tolerance studies at the NSLS and ESRF have used fruit-fly pupae<sup>21</sup>, rabbits, rats<sup>17, 18, 24-26</sup>, gerbils, mice<sup>14, 20, 22, 27</sup>, duck embryos<sup>12</sup>, and chick embryos<sup>10, 11</sup>.

### Conclusions

The 6 GeV ESRF ring is the only source of synchrotron radiation in Europe generating intense X ray microbeams for experimental MRT, having a broad energy spectrum of photons peaking in the 80 – 120 keV range and beam intensities high enough, potentially, to deliver an absorbed physical radiation dose to deep targets in large animals, small children, and adult humans; MRT requires the delivery of several hectogray doses within a fraction of a second, deep to the skin. Regulatory and logistic requirements for implementation of clinical MRT will be stringent. The impetus for investigating the potentially unique advantages of MRT for certain human diseases has been recognized and is sustained by the recent consensus of an ESRF scientific advisory panel of sufficient diversity and broad expertise in its membership to merit serious consideration by the ESRF directorate. Accordingly, we propose that ID17 be used to implement a large-animal veterinary MRT study for veterinary radio-oncology. In that way, a wider community of clinical veterinarians and physicians will be able to assess outcomes from MRT in relation to those from existing radiotherapies for similar lesions in large animals.

### References

- [1] Wagner HP. Cancer in childhood and supportive care. *Support Care Cancer* 1999; 7: 293-294.
- [2] Mulhern RK, Merchant TE, Gajjar A, Reddick WE, Kun LE. Late neurocognitive sequelae in survivors of brain tumours in childhood. *Lancet Oncol* 2004; 5: 399-408.
- [3] Ribi K, Relly C, Landolt MA, Alber FD, Boltshauser E, Grotzer MA. Outcome of medulloblastoma in children: long-term complications and quality of life. *Neuropediatrics* 2005; 36: 357-365.
- [4] Rutkowski S, Bode U, Deinlein F, Ottensmeier H, Warmuth-Metz M, Soerensen N, Graf N, Emser A, Pietsch T, Wolff JE, Kortmann RD, Kuehl J. Treatment of early childhood medulloblastoma by postoperative chemotherapy alone. *N Engl J Med* 2005; 352: 978-986.
- [5] Adler JR Jr, Colombo F, Heilbrun MP, Winston K. Toward an expanded view of radiosurgery. *Neurosurgery* 2004; 55: 1374-1376.
- [6] Köhler A. Une nouvelle méthode permettant de faire agir, dans la profondeur des tissus, de hautes doses de rayons Roentgen et un moyen nouveau de protection contre les radiodermites. *Annales d'Électrobiologie et de Radiologie* 1909; 10: 661-664.
- [7] Kondziolka D, Lunsford LD, Loeffler JS, Friedman WA. Radiosurgery and radiotherapy: observations and clarifications. *J Neurosurg* 2004; 101: 585-589.
- [8] Slatkin DN, Spanne P, Dilmanian FA, Sandborg M. Microbeam radiation therapy. *Med Phys* 1992; 19: 1395-1400.

- [9] Laissue J, Spanne PO, Dilmanian FA, Gebbers J-O, Slatkin DN: Zell- und Gewebeläsionen nach räumlich fraktionierter Mikro-Bestrahlung des ZNS mit Synchrotron-Photonen. *Schweiz Med Wochenschr* 1992; 122: 1627.
- [10] Blattmann H, Burkard W, Djonov V, Di Michiel M, Brauer E, Stepanek J, Bravin A, Gebbers JO, Laissue JA. Microbeam irradiation of the chorio-allantoic membrane (CAM) of chicken embryo. *Strahlentherapie und Onkologie* 2002; 178 (Suppl. June 1): 118
- [11] Blattmann H, Gebbers J-O, Bräuer-Krisch E, Bravin A, Le Duc G, Burkard W, Di Michiel M, Djonov V, Slatkin DN, Stepanek J, Laissue JA. Applications of synchrotron X-rays to radiotherapy. *Nucl Instr Meth Physics Res A* 2005; 548: 17-22.
- [12] Dilmanian FA, Morris GM, Le Duc G, Huang X, Ren B, Bacarian T, Allen JC, Kaled-Ezra J, Orion I, Rosen EM, Sandhu, T, Sathe P, Wu XY, Zhong Z, Shivaparasad HL. Response of avian embryonic brain to spatially segmented x-ray microbeams. *Cell Mol Biol* 2001; 47: 485-493.
- [13] Dilmanian FA, Button TM, Le Duc G, Zhong N, Peña LA, Smith JA, Martinez SR, Bacarian T, Tammam J, Ren B, Farmer PM, Kaled-Ezra J, Micca PL, Nawrocky MM, Niederer JA, Recksiek FP, Fuchs A, Rosen EM. Response of rat intracranial 9L gliosarcoma to microbeam radiation therapy. *Neuro-Oncol* 2002; 4: 26-38.
- [14] Dilmanian FA, Morris GM, Zhong N, Bacarian T, Hainfeld JF, Kaled-Ezra J, Brewington LJ, Tammam J, Rosen EM. (2003) Murine EMT-6 carcinoma: high therapeutic efficacy of microbeam radiation therapy. *Radiat Res* 159: 632-641.
- [15] Dilmanian FA, Qu Y, Liu S, Cool CD, Gilbert J, Hainfeld JF, Kruse CA, Laterra J, Lenihan D, Nawrocky MM, Pappas G, Sze C-I, Yuasa T, Zhong N, Zhong Z, McDonald JW. X-ray microbeams: Tumor therapy and central nervous system research. *Nucl Instr Meth Physics Res A* 2005; 548: 30-37.
- [16] Laissue JA, Spanne P, Dilmanian FA, Nawrocky MM, Gebbers J-O, Slatkin DN, Joel DD: Mikrobestrahlung von Gliosarkomen der Ratte: Zell- und Gewebeläsionen (Microbeam irradiation of rat gliosarcomas: Cell and tissue lesions). *Schweiz med Wochenschr* 1995; 125:1887.
- [17] Laissue JA, Geiser G, Spanne PO, Dilmanian FA, Gebbers JO, Geiser M, Wu XY, Makar MS, Micca PL, Nawrocky MM, Joel DD, Slatkin DN. Neuropathology of ablation of rat gliosarcomas and contiguous brain tissues using a microplanar beam of synchrotron-wiggler-generated X rays. *Int J Cancer* 1998; 78: 654-660.
- [18] Laissue JA, Lyubimova N, Wagner HP, Archer DW, Slatkin DN, Di Michiel M, Nemoz C, Renier M, Brauer E, Spanne PO, Gebbers JO, Dixon K, Blattmann H. Microbeam radiation therapy. *Proc SPIE* 1999; 3770: 38-45. .
- [19] Laissue JA, Blattmann H, Di Michiel M, Slatkin DN, Lyubimova N, Guzman R, Zimmermann W, Birrer S, Bley T, Kircher P, Stettler R, Fatzer R, Jaggy A, Smilowitz HM, Brauer E, Bravin A, Le Duc G, Nemoz C, Renier M, Thomlinson W, Stepanek J, Wagner HP. The weanling piglet cerebellum: a surrogate for tolerance to MRT (microbeam radiation therapy) in pediatric neuro-oncology. *Proc SPIE* 2001; 4508: 65-73.
- [20] Miura M, Blattmann H, Bräuer-Krisch E, Bravin A, Hanson AL, Nawrocky MM, Micca PL, Slatkin DN, Laissue JA. Radiosurgical palliation of aggressive murine SCCVII squamous cell carcinomas using synchrotron-generated X-ray microbeams. *Br J Radiol* 2006; 79: 71-75.
- [21] Schweizer PM, Spanne P, Di Michiel M, Jauch U, Blattmann H, Laissue JA: Tissue lesions caused by microplanar beams of synchrotron-generated x-rays in *Drosophila melanogaster*. *Int J Radiat Biol* 2000; 76 (4): 567-574.
- [22] Serduc R, Vérant P, Vial J-C, Farion R, Rocas L, Rémy C, Fadlallah T, Bräuer E, Bravin A, Laissue J, Blattmann H, van der Sanden B. In vivo two-photon microscopy study of short term effects of microbeam irradiation on normal mouse brain microvasculature . *Int J Radiat Oncol Biol Phys*; 2006; 64 (5):1519-1527.
- [23] Slatkin DN, Dilmanian FA, Nawrocky MM, Spanne P, Gebbers J-O, Archer DW, Laissue JA. Design of a multislit, variable width collimator for microplanar beam radiotherapy. *Rev Sci Instrum* 1995; 66:1459-1460.
- [24] Slatkin DN, Spanne P, Dilmanian FA, Gebbers JO, Laissue JA. Subacute neuropathological effects of microplanar beams of x-rays from a synchrotron wiggler. *Proc Natl Acad Sci USA* 1995; 92: 8783-8787. .
- [25] Smilowitz HM, Blattmann H, Bräuer-Krisch E, Bravin A, Di Michiel M, Gebbers J-O, Hanson AL, Lyubimova N, Slatkin DN, Stepanek J, Laissue JA. Synergy of gene-mediated immunoprophylaxis and microbeam radiation therapy (MRT) for advanced intracerebral rat 9L gliosarcomas. *J Neurooncol* 2006;78: 135-143.
- [26] Regnard P, Le Duc G, Bräuer-Krisch E, Troprès I, Siegbahn EA, Kusak A, Clair C, Bernard H, Dallery D, Laissue JA , Bravin A: Irradiation of intracerebral 9L gliosarcoma by a single array of microplanar

- X-ray beams from a synchrotron: balance between curing and sparing. *Phys Med Biol* 2008; 53: 861-878.
- [27] Serduc R, van de Looij Y, Francony G, Verdonck O, van der Sanden B, Laissue J, Farion R, Bräuer-Krisch E, Siegbahn EA, Bravin A, Prezado Y, Segebarth C, Rémy C, Lahrech H. Characterization and quantification of cerebral edema induced by synchrotron x-ray microbeam therapy. *Phys Med Biol* 2008; 53: 1153-1166.
- [28] Zhong N, Morris GM, Bacarian T, Rosen EM, Dilmanian FA. Response of rat skin to high-dose unidirectional x-ray microbeams: a histological study. *Radiat Res* 2003; 160:133-142.
- [29] Timmermann B, Kortmann RD, Kuhl J, Rutkowski S, Dieckmann K, Meisner C, Bamberg M. Role of radiotherapy in anaplastic ependymoma in children under age 3 years: results of the prospective German brain tumor trials HIT-SKK 87 and 92. *Radiother Oncol* 2005; 77: 278-285.
- [30] Laissue JA, Blattmann H, Wagner HP, MA Grotzer MA, Slatkin DN: Prospects for microbeam radiation therapy of children's brain tumours to reduce neurological sequelae. *Developmental Medicine and Child Neurology* 2007; 49: 577-581.